arXiv:hep-ph/0112009v1 2 Dec 2001

UCLA/01/TEP/34
February 1, 2008

Baryogenesis in the wake of inflation

Alexander Kusenko

Department of Physics and Astronomy, UCLA, Los Angeles, CA 90095-1547
and
RIKEN BNL Research Center, Brookhaven National Laboratory, Upton, NY 11973

Electroweak baryogenesis could be very efficient at the end of an electroweak-scale inflation. Reheating that followed inflation could create a highly non-equilibrium plasma, in which the baryon number violating transitions were rapid. In addition, the time-dependent motions of the scalar degrees of freedom could provide the requisite CP violation. If the final reheat temperature was below 100 GeV, there was no wash-out of the baryon asymmetry after thermalization. The observed value of the baryon asymmetry can be attained in a number of models, some of which do not require a significant departure from the Standard Model.

PRESENTED AT

COSMO-01
Rovaniemi, Finland,
August 29 – September 4, 2001

1 Introduction

Inflation probably took place in the early universe. If the scale of (the latest) inflation was of the order of the electroweak scale, the subsequent reheating would provide ideal conditions for electroweak baryogenesis.

The original scenario of Kuzmin, Rubakov, and Shaposhnikov [1, 2] was based on a brilliant idea that all three Sakharov's conditions [3] necessary for a successful baryogenesis were satisfied, at least qualitatively, in the early universe due to the properties of the Standard Model at finite temperature. First, the baryon number was violated by sphaleron transitions. Second, the universe was out of thermal equilibrium during a phase transition. Finally, CP was broken, because it is not an exact symmetry in the Standard Model.

Unfortunately, this scenario, in its original form, cannot account for the observed baryon-to-photon ratio $\eta \sim 10^{-10}$. First problem has to do with the fact that the electroweak phase transition is not sufficiently strongly first-order unless the Higgs mass is below about 45 GeV, which is ruled out by experiment. Second, the CP violation from the Cabibbo-Kobayashi-Maskawa matrix is way too small for baryogenesis because its contribution to η is suppressed by high powers of the Yukawa couplings.

At the end of inflation, however, the universe was emphatically out of equilibrium. Could the baryon number violating processes take place in that non-thermal environment? The answer is yes; this was clearly demonstrated by a consensus of numerical [6] and analytical [5, 6, 7] arguments. Finally, the requisite CP violation may come from time-dependent solutions for the scalar zero-modes in the background [7]. Altogether, electroweak baryogenesis can be very efficient at the end of inflation.

2 Electroweak baryogenesis at preheating

The idea [5, 6] of using for baryogenesis the non-equilibrium nature of preheating [9], along with the new sources of CP violation that preheating offers [7], is very appealing because it is simple and requires only a minor "upgrade" of the Standard Model. The inflaton has to be added to the Standard Model, but this is a widely accepted piece of new physics that is desirable for other reasons. The only uncomfortable features of the scenario proposed independently in Refs. [5] and [6] are the relatively low scale of inflation and the need to couple the inflaton to the Higgs sector. In combination with the constraint that inflation must last for more than 60 e-folds, this forces the inflaton potential to be unnaturally flat.

Of course, if the universe has undergone more than one period of inflation, the latest, electroweak-scale inflation could have lasted for only a small number of e-folds. In this case, there is no problem with naturalness.

Low-scale inflation models have been constructed [10]. One of the side benefits of lowering the inflation scale is avoiding the gravitino over-production constraints. There are other particle-physics motivations for using the TeV scale, which is associated with supersymmetry breaking in a class of models [11].

Although the simplest models of hybrid inflation do not yield parametric resonance [12], there is no fundamental reason why a weak-scale inflation could not be followed by resonant preheating [13]. The latter possibility allows for efficient electroweak baryogenesis at preheating.

2.1 Baryon number non-conservation during preheating



Figure 1: Evolution of the effective temperature with time in a numerical simulation of electroweak baryogenesis at preheating [6]

Numerical simulations in 1+1 dimensions [6, 14] demonstrate that the baryon number violating processes can go unsuppressed in a non-thermal plasma at preheating. One way to think about this is in terms of "effective temperature" [6]. Although plasma is far from equilibrium, the Chern-Simons number changes at the same rate as in thermal plasma at temperature T_{eff}. Since sphalerons, being extended objects, are mainly affected by the long-wavelength modes, it is the infra-red part of the spectrum that determines T_{eff}. At preheating, the long-wavelength modes are overpopulated as compared to the thermal ensemble. Hence, the effective temperature T_{eff} is higher than the energy density. When thermal equilibrium is restored, the final temperature $T_R < 100$ GeV $< T_{\text{eff}}$. At that point the rate of baryon number violating processes is negligible. This behavior, favorable to baryogenesis, is shown in Figs. 1,2.

One can try to describe this behavior semi-analytically by isolating certain gauge degrees of freedom and studying their evolution at preheating [7], when the vector boson mass, M_W changes with time due to the oscillations of the inflaton coupled to the Higgs sector. Let us introduce an *ansatz* [15, 16] for the gauge potential

$$gA_\mu = (\frac{\tau_a}{2i})A^a_\mu. \tag{1}$$



Figure 2: Change in Chern-Simons number during the run shown in Fig. 1

Our spatially-homogeneous *ansatz* is:

$$gA_0 = 0;\ gA_i = (\frac{\tau_i}{2i})\phi(t) \tag{2}$$

in which the group index is tied to the spatial index. By the conventional rules of charge conjugation and parity for the gauge potential, ϕ is C even, P odd, CP odd.

It is important to note that this *ansatz* does not correspond to a non-vanishing VEV for an EW field. Gauge invariance alone is enough to ensure that there can be no expectation value coupling the space-time indices to group indices.

One readily calculates the EW electric and magnetic fields:

$$gE_i \equiv G_{0i} = (\frac{\tau_i}{2i})\dot{\phi}(t);\ gB_i \equiv \frac{1}{2}\epsilon_{ijk}G_{jk} = (\frac{\tau_i}{2i})\phi^2. \tag{3}$$

Then one calculates the density W of the Chern-Simons number as:

$$W = (\frac{1}{8\pi^2})\phi^3. \tag{4}$$

It is straightforward to check that $\dot{W}$ is the topological charge density Q, related to B+L violation through the anomaly equation.

With the assumption of a given Higgs VEV, the equations of motion for the gauge potential are:

$$[D^\mu, G_{\mu\nu}] + M_W^2(t)(A_\nu + (\partial_\nu U)U^{-1}) = 0. \tag{5}$$

Here the unitary matrix U represents the Goldstone (phase) part of the Higgs field. The mass term will be assumed to have the form:

$$M_W^2(t) = m^2(1 + \epsilon\cos(\omega t)) \tag{6}$$

where m is the value of M_W with no oscillations.

This *ansatz* has some important properties. First, it carries some non-zero Chern-Simons density. Second, the equation of motion for the field ϕ,

$$\ddot{\phi} + 2\phi^3 + (1 + \epsilon \cos rt)\phi = 0, \tag{7}$$

has resonantly growing solutions. Finally, the ansatz can evolve for a long time without breaking spatial homogeneity [7].



Figure 3: Resonant growth of the Chern-Simons number density in the gauge "condensate" for the case of slowly changing driving frequency [7].

The Chern-Simons density grows as the third power of ϕ.

2.2 CP violation at preheating

CP and time-reversal symmetries at preheating may be broken by the time-dependent classical motions of scalar fields in the background [8]. This is similar to the source of CP violation used by Affleck and Dine [17]. CP violation of this kind is poorly constrained by experiment. Several viable scenarios for electroweak baryogenesis at preheating were presented in Ref. [8]. For example, a modified spontaneous baryogenesis a la Cohen, Kaplan, and Nelson [18] becomes very efficient at preheating. Their original scenario used the variation of the Higgs field inside a wall of a bubble formed in a first-order phase transition. A similar effect can occur at preheating uniformly in space, on the horizon scales [8]. One can get the desired baryon asymmetry in a Standard Model supplemented by an additional Higgs doublet and an inflaton sector [8]. The difference with the scenario proposed by Cohen, Kaplan and Nelson [18] is that in our case CP violation occurs homogeneously in space, as opposed to in a bubble wall. In addition, the final prediction for baryon asymmetry in the CKN scenario was very far from the equilibrium value because the sphaleron rate was slow on the time scales associated with the growth of the bubble. In our case, the Higgs parameters change slowly in time while the baryon number non-conservation is rapid. This allows a slow adiabatic adjustment of the baryon number to that which minimizes the free energy.

3 Other possibilities for baryogenesis at preheating

Non-thermal production of heavy particles at the end of inflation [19] could result in baryogenesis [20], or a leptogenesis [21] via the decay of the heavy particles.

Alternatively, in models with low-energy supersymmetry, weak-scale inflation can lead to formation of the Affleck-Dine condensate [17, 22]. In general, this condensate does not remain homogeneous and can break up [23] into SUSY Q-balls [24]. This affords a number of interesting possibilities for generating baryons and dark matter simultaneously [23, 25, 26, 27].

If there were complex scalar fields not associated with supersymmetry, their fragmentation in a similar fashion could produce stable Q-balls that could make up the dark matter. Interestingly enough, this form of dark matter can be strongly self-interacting [28, 29] and can, therefore, resolve the problems with non-interacting cold dark matter on small scales [30].

4 Conclusion

A highly non-thermal state of the universe at the end of inflation provides a fertile ground for baryogenesis. In addition to a dramatic departure from thermal equilibrium, preheating creates conditions for baryon number non-conservation and activates sources of CP violation that are largely unconstrained by the experimental data. Several modest modifications of the Standard Model allow for efficient electroweak baryogenesis in the wake of inflation.

References

[1] V. A. Kuzmin, V. A. Rubakov, and M. Shaposhnikov, Phys. Lett. **155B**, 36 (1985).

[2] V. A. Rubakov and M. E. Shaposhnikov, Usp. Fiz. Nauk **166**, 493 (1996) [Phys. Usp. **39**, 461 (1996)];

[3] A. D. Sakharov, JETP Lett. **6**, 24 (1967).

[4] V. A. Kuzmin, Pisma ZhETP, **13**, 335 (1970).

[5] L. M. Krauss and M. Trodden, Phys. Rev. Lett.**83**, 1502 (1999).

[6] J. Garcı́a-Bellido, D. Grigoriev, A. Kusenko, and M. Shaposhnikov, Phys. Rev. D **60**, 123504 (1999).

[7] J. M. Cornwall and A. Kusenko, Phys. Rev. D **61**, 103510 (2000)

[8] J. M. Cornwall, D. Grigoriev and A. Kusenko, Phys. Rev. D **64**, 123518 (2001)

[9] L. Kofman, A. Linde, and A. A. Starobinsky, Phys. Rev. Lett. **73**, 3195 (1994), *ibid.* **76**, 1011 (1996); Phys. Rev. D **56**, 3258 (1997).

[10] A.D. Linde, Phys. Lett. **B259**, 38 (1991); Phys. Rev. D **49**, 748 (1994); L. Knox and M. Turner, Phys. Rev. Lett. **70** (1993) 371; L. Randall, M. Soljačić, and A. H. Guth, Nucl. Phys. B **472**, 377 (1996); J. García-Bellido, A. D. Linde and D. Wands, Phys. Rev. D **54**, 6040 (1996); D. H. Lyth and E. D. Stewart, Phys. Rev. D **54**, 7186 (1996); J. García-Bellido, these Proceedings.

[11] G. F. Giudice and R. Rattazzi, Phys. Rept. **322**, 419 (1999).

[12] G. N. Felder, J. Garcia-Bellido, P. B. Greene, L. Kofman, A. D. Linde and I. Tkachev, Phys. Rev. Lett. **87**, 011601 (2001).

[13] A. Linde, private communications.

[14] J. Garcia-Bellido and D. Y. Grigoriev, JHEP **0001**, 017 (2000). A. Rajantie, P. M. Saffin and E. J. Copeland, arXiv:hep-ph/0010347; E. J. Copeland, D. Lyth, A. Rajantie and M. Trodden, Phys. Rev. D **64**, 043506 (2001); A. Rajantie, these Proceedings, hep-ph/0111200.

[15] K. M. Bitar and S. Chang, Phys. Rev. **D17**, 486 (1978).

[16] J. M. Cornwall, Phys. Rev. **D40**, 4130 (1989).

[17] I. Affleck and M. Dine, Nucl. Phys. B **249**, 361 (1985).

[18] A. G. Cohen, D. B. Kaplan and A. E. Nelson, Phys. Lett. B **263**, 86 (1991).

[19] D.J. Chung, E.W. Kolb, and A. Riotto, Phys. Rev. Lett. **81**, 4048 (1998); Phys. Rev. **D59**, 023501 (1999).

[20] E. W. Kolb, A. D. Linde and A. Riotto, Phys. Rev. Lett. **77**, 4290 (1996).

[21] G. F. Giudice, M. Peloso, A. Riotto and I. Tkachev, JHEP **9908**, 014 (1999).

[22] M. Dine, L. Randall and S. Thomas, Nucl. Phys. **B458** (1996) 291; A. Anisimov and M. Dine, arXiv:hep-ph/0008058.

[23] A. Kusenko, M. Shaposhnikov: Phys. Lett. B **418**, 46 (1998)

[24] A. Kusenko, Phys. Lett. B **405**, 108 (1997); Phys. Lett. B **404**, 285 (1997); A. Kusenko, V. Kuzmin, M. E. Shaposhnikov and P. G. Tinyakov, Phys. Rev. Lett. **80**, 3185 (1998).

[25] K. Enqvist, J. McDonald: Phys. Lett. B **425**, 309 (1998); Nucl. Phys. B **538**, 321 (1999); Phys. Rev. Lett. **81**, 3071 (1998); Phys. Lett. B **440**, 59 (1998); Phys. Rev. Lett. **83**, 2510 (1999)

[26] S. Kasuya and M. Kawasaki, Phys. Rev. D **61**, 041301 (2000); Phys. Rev. D **62**, 023512 (2000); Phys. Rev. D **64**, 123515 (2001) [arXiv:hep-ph/0106119]. S. Kasuya, Phys. Lett. B **515**, 121 (2001) [arXiv:astro-ph/0105408].

[27] M. Laine and M. E. Shaposhnikov, Nucl. Phys. B **532**, 376 (1998) [arXiv:hep-ph/9804237].

[28] A. Kusenko and P. J. Steinhardt, Phys. Rev. Lett. **87**, 141301 (2001) [arXiv:astro-ph/0106008].

[29] K. Enqvist, A. Jokinen, T. Multamaki and I. Vilja, arXiv:hep-ph/0111348.

[30] D. N. Spergel and P. J. Steinhardt, Phys. Rev. Lett. **84**, 3760 (2000); P. Steinhardt, these Proceedings.